


Acute Chest Pain Pathway



*Avoid fondaparinux if CrCl<20ml/min – can use appropriately dosed low molecular weight heparin as alternative (refer to [ADTC 176](#)).
 ** Patients who receive PCI will be switched to prasugrel (by interventional centre) unless contraindicated (previous TIA, CVA, ICH / patients on anticoagulation / severe hepatic impairment / patients for thrombolysis / propensity to bleeding (anaemia, GI bleed)).
 ***Please note results for POC Trop I and Lab Trop T are not interchangeable or directly comparable

Appendix 1

GJNH High Risk NSTEMI pathway

HEART



History (Anamnesis)	Highly suspicious	2	
	Moderately suspicious	1	
	Slightly suspicious	0	
ECG	Significant ST-deviation	2	
	Non-specific repolarisation disturbance/LBBB/PM	1	
	Normal	0	
Age	≥ 65 years old	2	
	45-64 years old	1	
	≤ 45 years old	0	
Risk Factors	≥ 3 risk factors or history of atherosclerotic disease	2	
	1 or 2 risk factors	1	
	No known risk factors	0	
Troponin	≥ 29 ng/l	2	
	5 – 28 ng/l	1	
	< 5 ng/l	0	
		Total	

Risk Factors:

- Hypercholesterolaemia
- Hypertension
- Diabetes Mellitus
- Cigarette smoking
- Positive family history
- Obesity (BMI>30)

***HOW TO CALCULATE THE HEART SCORE**

The HEART score is a risk stratification tool first used in the Emergency Department to predict the likelihood of a major adverse cardiac event within 6 weeks following presentation with chest pain.

A score is assigned from 5 specific elements (History, ECG changes, Age, Risk factors and Troponin) to give a value between 0 and 9. Three of the elements are explained in detail below:

History - From your history characterise the patient's chest pain as typical or atypical. The following distinctions have been agreed:

1. **Typical pain** - central of left-sided chest pain with radiation to the arms or throat, or associated sweating or clamminess.

2. **Atypical pain** - without chest pain or right sided chest pain or pain that radiates to the back or is worsened by inspiration/palpation.

→ 2 points: highly suspicious chest pain (i.e. typical pain)

→ 1 point: moderately suspicious chest pain (i.e. mixed typical/atypical features)

→ 0 point: chest pain slightly or moderately suspicious

Electrocardiogram (12 Lead ECG) - From the 12 lead ECG:

→ 2 points: ECG shows features new/presumed new features of acute ischaemia or infarction (eg. significant ST depression, T-wave inversion)

→ 1 point: ECG is abnormal but not diagnostic of ischaemia (eg. right bundle branch block, paced rhythm) or if ECG suggests previous infarction

→ 0 points: ECG is normal

Risk Factors: Count the number of risk factors for coronary artery disease:

Diabetes mellitus

Current or recent (<90 days) smoker

Hypertension (diagnosed or treated)

Hypercholesterolaemia

Family history of coronary disease


Obesity (BMS >30)

→ 2 points: 3 or more risk factors or significant atherosclerotic disease (including previous coronary revascularisation, myocardial infarction, peripheral arterial disease)

→ 1 point: 1-2 risk factors

→ 0 point: no risk factors

HEART score reliably predicts endpoints



HEART	~ % pts	MACE/n	MACE	Death	Proposed Policy
0-3	32%	38/1993	1.9%	0.05%	Discharge
4-6	51%	413/3136	13%	1.3%	Observation, risk management
7-10	17%	518/1045	50%	2.8%	Observation, treatment, CAG

*MACE = Major Adverse Cardiac Event = Myocardial Infarction, PCI/CABG, all-cause death. Based on N=6174

<http://www.heartscore.nl/>

Appendix 2

Diagnosis of Myocardial Infarction with High Sensitivity Cardiac Troponin (hs-cTn)

Elevated hs-cTn level should always be interpreted in the context of patient symptoms and ECG findings. Typical chest pain with ECG changes makes * Type 1 MI likely. Conversely, in the absence of evidence of ischaemia, other causes should be sought. Please note that results for Troponin I (POC trop) and Troponin T (Lab trop) are not directly interchangeable. This must be taken into account when assessing changes in Troponin level over 6 hours.

Causes of Elevated hs-cTn other than *Type 1 MI (*MI due to a primary coronary event, usually atherosclerotic plaque rupture)

Acute Conditions	Chronic Conditions
Imbalance of Demand/Supply (Type 2 MI)	
Tachy- or bradyarrhythmias	Tachy- or bradyarrhythmias
Aortic Dissection	
	Severe aortic valve stenosis
Cardiogenic, hypovolaemic and septic shocks	
	Anaemia
	Hypertension
	Left Ventricular Hypertrophy
Coronary Embolism or Vasculitis	
Coronary Spasm	Coronary Spasm
Endothelial dysfunction	Endothelial dysfunction
Cocaine use	
Non-ischaemic Myocardial Damage	
Cardiac contusion	
Cardiac surgery	
Radiofrequency or cryoblation therapy	
Pacing or defibrillation shocks	Pacing or defibrillation shocks
Rhabdomyolysis with cardiac involvement	
Myopericarditis	Myopericarditis
Cardiotoxic agents	Cardiotoxic agents
Some chemotherapeutics	Some chemotherapeutics
Carbon monoxide poisoning	
Multifactorial Causes of Myocardial Damage	
Heart Failure	Heart Failure
Takotsubo cardiomyopathy	
Severe pulmonary embolism	
	Pulmonary Hypertension
Extreme exertion	
Sepsis	
Gastrointestinal bleeding	
Rhabdomyolysis without cardiac involvement	
Renal Failure	
	Infiltrative diseases such as sarcoidosis or amyloidosis
Severe acute neurological disease, such as stroke or trauma	
Skeletal myopathies	

2023 ESC Guidelines for the management of acute coronary syndromes: Developed by the task force on the management of acute coronary syndromes of the European Society of Cardiology (ESC)

<https://doi.org/10.1093/eurheartj/ehad191>